

Parametric Decay of the Curvaton

K. Enqvist and S. Nurmi

*Department of Physics, University of Helsinki, and Helsinki Institute of Physics,
P.O.Box 64, FIN-00014, University of Helsinki, Finland*

G. I. Rigopoulos

Helsinki Institute of Physics, P.O.Box 64, FIN-00014, University of Helsinki, Finland

We argue that the curvaton decay takes place most naturally by way of a broad parametric resonance. The mechanism is analogous to resonant inflaton decay but does not require any tuning of the curvaton coupling strength to other scalar fields. For low scale inflation and a correspondingly low mass scale for the curvaton, we speculate on observable consequences including the possibility of stochastic gravitational waves

INTRODUCTION

The curvaton scenario [1–4] provides an alternative mechanism for producing the primordial density perturbations. The curvaton is a light field which is subdominant during inflation but whose energy density will eventually become an important (if not actually the dominant) energy component in the universe. Because of its lightness, during inflation the curvaton acquires fluctuations which at the time of curvaton decay are transformed into curvature perturbations.

So far, the decay of the curvaton has mostly been treated as a perturbative process, see e.g. [5, 6], where the curvaton condensate decays into relativistic particles, determined by a rate Γ and occurring roughly when $H \sim \Gamma$. This happens at the stage when the curvaton field is oscillating about the minimum of its potential, whence it behaves like pressureless dust so that its relative energy density grows with respect to radiation produced by the inflaton decay. The curvature perturbation induced by the curvaton decay is mostly adiabatic but can give rise to a large non-gaussianity [6, 7] if the curvaton energy density is too small¹; this implies a lower limit on the fraction of the energy density in the curvaton condensate at the decay time. Another generic feature of curvaton models with no mixed inflaton and curvaton perturbations is the small amplitude tensor perturbations [2, 6]; this is a consequence of the low inflationary scale required to keep the inflaton perturbations negligible.²

However, since the curvaton is a coherently oscillating condensate, it is possible to excite other boson fields in a regime of parametric resonance, in a manner similar to inflationary preheating [10, 11]. As a consequence, the curvaton decay process could have important cosmological ramifications for e.g. non-gaussianities, which are known to be enhanced [12, 13] in some preheating scenarios, as well as for stochastic gravitational waves, as will be discussed below. Preheating is a violent process in which the occupation numbers of fields coupled to the curvaton grow exponentially. In effect, dense lumps of decay products are created very rapidly and randomly in the real space, and as these highly inhomogeneous concentrates of energy density collide, there arise large-amplitude, high-frequency gravitational waves, as has been discussed recently [14, 15]. These do not leave a signal on the CMB but could possibly be observed in current or future gravitational wave detectors such as LIGO [16], LISA [17] or BBO [18].

In the present paper we study the possibility of preheating in generic curvaton models and explore the main consequences of such a scenario (preheating has been discussed in the context of MSSM flat direction curvatons in [19]). Indeed, the curvaton has to couple to other fields if it is to decay; if it couples to other scalars, then there would be two options. Either the additional scalars would be light and shifted from their minima during inflation, in which case they would simply be additional curvatons, or they could be massive and driven to their minima. In the first case one still would have to find a way for the curvaton(s) to decay. However, in the latter case there can arise a parametric resonance during curvaton oscillations with the ensuing rapid decay of the curvaton.

We find that the preheating process in the curvaton scenario is qualitatively similar to the preheating after inflation although the resonance dynamics can be somewhat altered. This is because the curvaton, unlike the inflaton, could remain subdominant during the entire preheating stage. Moreover, in the scenario studied here the backreaction of the excited boson fields always becomes significant at the end of preheating whereas in the standard inflationary

¹ Strictly speaking, this holds only for a quadratic curvaton potential, see [8].

² See however [9] where it is demonstrated that the amplitude of CMB-scale tensor perturbations in the curvaton model can be directly proportional to the amount of the produced non-gaussianity due to second order effects.

case the preheating may terminate already at an earlier stage. Since the gaussian part of curvaton perturbations on superhorizon scales is not affected by preheating and the final decay of the curvaton will eventually take place in a perturbative reheating process, the possible preheating stage does not alter the standard predictions of the curvaton model [6]. However, interestingly enough we find that preheating may generate stochastic gravitational waves with an amplitude much larger than the inflationary tensor perturbations in the curvaton model, and we discuss the feasibility of observing such signatures of curvaton preheating. The study of the non-gaussianities is left for a future work.

THE INFLATIONARY PERIOD

Preheating, or resonant decay of the curvaton, may occur in a large class of curvaton models with several interacting scalar fields, as we will now discuss. To illustrate the main features of the process, let us consider a very simple model with the potential

$$V = V(\phi) + V(\sigma, \chi) , \quad (1)$$

where ϕ is the inflaton, σ is the curvaton and χ is an additional scalar field which the curvaton decays into. Note that χ should not be yet another curvaton. We assume that the inflationary dynamics is completely determined by the inflaton field, $H_* = H_*(\phi)$, which requires that the energy densities of the curvaton and the χ field are much smaller than $H_*^2 M_{\text{P}}^2$. We also assume that all fields have canonical kinetic terms.

Let us further assume that

$$V(\sigma, \chi) = \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} g^2 \sigma^2 \chi^2 + \frac{1}{4!} \lambda \chi^4 , \quad (2)$$

where $g \lesssim 1$ and $\lambda \lesssim 1$ are some coupling constants. For certain field and coupling values, both σ and χ could act as the curvaton but this is not the case we wish to study here. We therefore concentrate on the regime where the effective mass of the χ field is larger than the Hubble parameter $g\sigma \gtrsim H_*$ and the contribution to the curvaton mass coming from the χ field can be neglected $\chi \ll m_\sigma/g$. In this case the field χ will not acquire perturbations during inflation and it simply represents a generic bose field that is coupled to curvaton but will not itself act as a curvaton. Under these assumptions, the value of the χ field is driven towards zero exponentially fast during inflation.

The curvaton is effectively massless during inflation $m_\sigma \ll H_*$ and acquires perturbations proportional to the Hubble parameter $\delta\sigma \sim H_*$. In the standard scenario where the curvaton is assumed to decay perturbatively at some stage after inflation, the amplitude of primordial curvature perturbations can be estimated by $\mathcal{P}_\zeta^{1/2} \sim r H_*/\sigma$ [2, 6], where $r \sim \rho_\sigma/\rho$ measures the fraction of the curvaton of the total energy density at the decay time. Since the curvaton is very light, its equation of motion is approximatively given by $3H_*\dot{\sigma} \simeq -m_\sigma^2 \sigma$ (for a discussion of curvaton dynamics in various potentials, see [20]). If inflation lasts long enough, the curvaton will eventually roll down to the region in its potential where the quantum effects overcome the classical drift, $|\dot{\sigma}| \lesssim H_*^2$. The typical curvaton value during inflation in this case is thus given by

$$\sigma_* \sim H_* \left(\frac{H_*}{m_\sigma} \right)^2 . \quad (3)$$

In the limit $r \ll 1$, this yields an estimate

$$r_{\text{osc}} \sim \left(\frac{m_\sigma}{M_{\text{P}}} \right)^2 \left(\frac{H_*}{m_\sigma} \right)^6 \quad (4)$$

for the curvaton contribution to the total energy density at the beginning of oscillations.

THE PARAMETRIC RESONANCE

Preheating in the curvaton model can take place when the curvaton starts to oscillate after inflation and gives a time dependent effective mass for the χ field that initially has a vanishing vev. The situation is analogous to the standard case of preheating in the conventional inflationary models [10, 11]. As usual, an efficient preheating requires broad resonance bands which implies the condition

$$q \equiv \frac{g^2 \sigma^2}{4m_\sigma^2} \gg 1 \quad (5)$$

at the beginning of oscillations. In our case, this necessarily holds as a consequence of the mass condition for the χ field, $g \gtrsim H_*/\sigma$, combined with the lightness of the curvaton $m_\sigma \ll H_*$.

Depending on the fraction of the energy density in the curvaton field at the time of oscillations, denoted as r_{osc} , there are two different possibilities for realizing the preheating stage. If the curvaton dominates the energy density at the beginning of oscillations with $r_{\text{osc}} \sim 1$, preheating proceeds exactly in the same manner as in the standard inflationary case. In the opposite limit, $r_{\text{osc}} \ll 1$, the universe is not dominated by the oscillating curvaton but by the inflaton decay products assumed to be homogeneous radiation. This however leads only to minor differences as compared to the standard case [11] as we will discuss below.

In the limit $r_{\text{osc}} \sim 1$ the universe is effectively matter dominated during the curvaton oscillations. The time evolution of the curvaton is given by

$$\sigma(t) = \frac{\sigma_*}{m_\sigma t} \sin(m_\sigma t) \equiv \bar{\sigma} \sin(m_\sigma t) , \quad (6)$$

where $\bar{\sigma} \propto t^{-1}$ measures the amplitude of oscillations. For definiteness, we choose $t_0 = \pi/(2m_\sigma)$ as the beginning of oscillations and set $a_0 = 1$. The initial amplitude is then given by $\bar{\sigma} = 2\sigma_*/\pi$, where σ_* is the value during inflation. Using a rescaled field variable $X_k = a^{3/2}\chi_k$, the equation of motion for the χ field in Fourier space reads

$$\ddot{X}_k + \left(\frac{k^2}{a^2} + g^2 \sigma^2 \right) X_k = 0 , \quad (7)$$

where terms proportional to $H^2 \ll g^2 \sigma^2$ and the small contribution from the quartic term in (2) have been neglected. The solutions can be formally written as

$$X_k(t) = \frac{\alpha_k(t)}{\sqrt{2\omega}} e^{-i \int^t \omega dt} + \frac{\beta_k(t)}{\sqrt{2\omega}} e^{i \int^t \omega dt} , \quad (8)$$

and the occupation number of a given mode is determined by $n_k = |\beta_k|^2$. In the broad resonance regime, the occupation numbers n_k remain constant for most of the time since the mass of the χ field is slowly varying and much larger than the curvaton mass m_σ [11]. However, as the oscillating curvaton crosses the origin during each cycle of oscillation, the χ field becomes effectively massless and its frequency $\omega_k = \sqrt{(k/a)^2 + g^2 \sigma^2}$ changes in a non-adiabatic fashion $\dot{\omega}_k \gtrsim \omega_k^2$. This can lead to copious production of χ particles via a resonant decay of the curvaton condensate.

The particle production takes place at very short time intervals around times t_n defined by $\sigma(t_n) = 0$. The curvaton value in the vicinity of these intervals can be approximated by $|\sigma| \simeq \bar{\sigma}_n m_\sigma (t - t_n)$. By substituting this into (7), the equation of motion becomes

$$\frac{d^2 X_k}{d\tau^2} + (\kappa_n^2 + \tau_n^2) X_k = 0 , \quad (9)$$

where we have introduced the variables $k_* \equiv \sqrt{gm_\sigma \bar{\sigma}}$, $\kappa_n = k/(ak_*)$, $\tau_n = k_*(t - t_n)$ and all the time arguments not explicitly written out are understood to be evaluated at t_n . The solutions for this equation are parabolic cylinder functions $W(-\kappa_n^2/2; \pm\sqrt{2}\tau_n)$. By comparing them in the subsequent adiabatic regimes $t_{n-1} < t < t_n$ and $t_n < t < t_{n+1}$ one can estimate the change of occupation numbers n_k around the times t_n when the system is non-adiabatic. In the limit of large occupation numbers, $n_k \gg 1$, the result is given by [11]

$$n_k^{n+1} = n_k^n \exp(2\pi\mu_k^n) , \quad (10)$$

where $2\pi\mu_k^n = \ln(1 + 2e^{-\pi\kappa_n^2} - 2\sin\theta_n e^{-\pi\kappa_n^2/2} \sqrt{1 + e^{-\pi\kappa_n^2}})$ and θ_n is a phase factor that is essentially a random variable for each resonance time t_n . This is why preheating in an expanding space has a stochastic nature.

The system formally describes scattering of waves in a parabolic potential. Each resonance time t_n , when the curvaton hits the origin, corresponds to a scattering event and changes the occupation numbers of the χ modes. The changes can be significant in the regime $\pi\kappa_n^2 \lesssim 1$ which corresponds to the momentum band

$$\frac{k^2}{a^2(t_n)} \lesssim \frac{k_*^2(t_n)}{\pi} = \frac{gm_\sigma \bar{\sigma}_n}{\pi} . \quad (11)$$

On average, the occupation numbers grow exponentially in this band, although the exponent μ_k in (10) can also be negative for some scattering events t_n , and the comoving particle number density as a function of time can be estimated as

$$n(t) = \frac{1}{(2\pi a)^3} \int d^3k n_k(t) \sim 10^{-3} \frac{k_*^3}{a^3 \sqrt{\mu m_\sigma t}} e^{2\mu m_\sigma t} . \quad (12)$$

Here μ is the maximum of the effective growth index μ_k^{eff} obtained from equation (10) by defining $\mu_k^{\text{eff}} t = \int^t dt \mu_k(t)$.

As the modes of the χ field grow exponentially, the term $g^2 \chi^2 \sigma^2$ in the curvaton potential becomes comparable to the mass term $m_\sigma^2 \sigma^2$ and the backreaction effects which eventually terminate the resonance need to be taken into account. The backreaction becomes important [11] when the number density of χ particles (12) grows to $n(t_{\text{br}}) \sim m_\sigma^2 \bar{\sigma}/g$ which yields an equation

$$t_{\text{br}} \sim \frac{1}{4\mu m_\sigma} \ln \left(\frac{10^6 \mu m_\sigma^4 t_{\text{br}}^3}{g^5 \sigma_*} \right), \quad (13)$$

from which t_{br} can be solved. The condition for broad resonance (5) becomes violated when $g\bar{\sigma}(t_f) \sim m_\sigma$, which yields

$$t_f \sim \frac{g\sigma_*}{m_\sigma^2} \gtrsim \frac{H_*}{m_\sigma^2}. \quad (14)$$

The last inequality follows from the condition $g \gtrsim H/\sigma$ which is required to keep χ massive during inflation. The backreaction generically takes place much before the end of broad resonance since the above equations yield $t_{\text{br}} \ll t_f$ for $m_\sigma/H_* \lesssim 10^{-2}$, assuming the curvaton value during inflation is given by equation (3). The interaction between the curvaton condensate and the χ particles breaks the condensate into curvaton particles which eventually come to dominate over the χ particles [11]. As usual, preheating needs to be followed by a reheating stage which eventually yields the hot big bang era. The simplest assumption [11] is that preheating just prepares slightly different initial conditions for the reheating stage during which the curvaton decays into standard model fields. In this case the standard predictions of the curvaton scenario for the gaussian part of density fluctuations remain largely unaltered. It is possible though that the non-thermal epoch after preheating could significantly affect the picture and lead to serious constraints on the curvaton scenario; this question should be addressed numerically by studying the thermalization process. However, even if the gaussian density fluctuations would not be affected, curvaton preheating leads to production of gravitational waves which could be detectable as discussed below. In addition, although we do not discuss this possibility further here, we note that preheating could also generate a significant level of non-gaussian density perturbations as can happen in the usual inflationary case [12, 13].

If the curvaton is subdominant at the beginning of oscillations, $r_{\text{osc}} \ll 1$, the above discussion gets slightly modified but the qualitative results remain the same. The exact time evolution of the curvaton in the radiation dominated background is given by Bessel functions but the solution is well approximated by

$$\sigma(t) \simeq \frac{\sigma_*}{(m_\sigma t + \frac{\pi}{8})^{3/4}} \sin \left(m_\sigma t + \frac{\pi}{8} \right). \quad (15)$$

This differs from the matter dominated case (6) only by a phase and the slightly slower decrease of the amplitude $\bar{\sigma} \propto t^{-3/4}$. It is therefore rather evident that the analysis of preheating will not significantly differ for the matter dominated case. The equation of motion for the χ field can be expressed exactly in the same form as equation (9) except that the time evolution of the variables κ_n and τ_n differs from the matter dominated case due to the different evolution of the scale factor $a(t)$ and the curvaton amplitude $\bar{\sigma}$. This does not affect the stochastic nature of the successive scatterings at times t_n and the occupation numbers of the χ field grow essentially at the same rate as before. We therefore expect that the number density of χ particles in the $r_{\text{osc}} \ll 1$ limit can be estimated by the same equation (12) as in the case $r_{\text{osc}} \sim 1$. This conclusion can be checked by solving the equations of motion for the curvaton and the χ field numerically. As illustrated in Fig. (1), the results in the matter and radiation dominated cases indeed coincide quite precisely in the regime where the backreaction of χ particles can be neglected. The time when the number density of the χ particles has grown large enough to yield a significant backreaction can be estimated as before using equation (12) and the condition $n(t_{\text{br}}) \sim m_\sigma^2 \bar{\sigma}/g$. Due to the smaller suppression by the scale factor in (12), the backreaction will occur slightly earlier than in the matter dominated case, as also seen in Fig. (1), but the effect is very small. In analyzing the dynamics after the onset of backreaction, the expansion of the universe can be neglected [11] and the radiation domination should not affect the situation here either.

GRAVITY WAVES

Since the decay of the curvaton can take place in a regime of parametric resonance, as was demonstrated above, it is natural to expect that gravitational waves will also be produced when it occurs. Indeed, the physics is similar as in the inflationary preheating: an oscillating field excites the production of particles in other bose fields at the times




FIG. 1: Comparison of resonant decay of the curvaton in radiation and matter dominated backgrounds. The figure shows $\ln n$ for the χ field in the cases $r=1$ (dashed lines) as well as $r < 1$ (solid lines). It also shows the limits above which backreaction becomes important in both cases. The parameters for these plots are $(M_P = 1)$, $m = 10^{-10}$, $g = 10^{-4}$, and $\sigma_* = 0.0002$. The onset of backreaction takes place after about 30 - 35 oscillations of the curvaton which corresponds to growth index $\mu \simeq 0.07$.

when their effective frequencies change non-adiabatically. According to [15], in such circumstances the generation of gravitational waves is inevitable. In what follows we make extensive use of the analysis in [15] which can be directly applied to our case with appropriate modifications.

Let us first reiterate the salient features of production of stochastic gravitational waves, which can be represented as the transverse-traceless part of the metric perturbations

$$ds^2 = a^2(\eta) [-d\eta^2 + (\delta_{ij} + h_{ij}) dx^i dx^j], \quad (16)$$

where $\partial_i h_{ij} = h_{ii} = 0$ and $d\eta = a^{-1}dt$. Gravitational waves are sourced by the transverse-traceless part of the energy momentum tensor T_{ij}^{TT} ³

$$\partial_\eta^2 \bar{h}_{ij}(\mathbf{k}) + \left(k^2 - \frac{\partial_\eta^2 a}{a} \right) \bar{h}_{ij}(\mathbf{k}) = 16\pi G a(\eta) T_{ij}^{TT}(\mathbf{k}), \quad (17)$$

where $\bar{h}_{ij} = ah_{ij}$. In the case of radiation domination $\partial_\eta^2 a = 0$ so equation (17) can be solved with the appropriate Green functions of the harmonic oscillator. Since we will be considering modes with $k \gg aH$ the same applies to a good approximation for the matter dominated case. The energy density in gravitational waves for subhorizon modes can then be written as [15]

$$\rho_{\text{gw}} = \frac{1}{32\pi G a^4} \langle \partial_\eta \bar{h}_{ij} \partial_\eta \bar{h}_{ij} \rangle. \quad (18)$$

Considering the gravitational waves being produced up to a time η_f we have

$$\left(\frac{d\rho_{\text{gw}}}{d \ln k} \right)_{\eta > \eta_f} = \frac{S_k(\eta_f)}{a^4(\eta)}, \quad (19)$$

where S_k for a universe filled with a scalar field χ reads

$$S_k(\eta_f) = \frac{2G}{\pi} k^3 \int \frac{d^3 \mathbf{p}}{(2\pi)^3} p^4 \sin^4(\hat{\mathbf{k}}, \hat{\mathbf{p}}) \left\{ \left| \int_0^{\eta_f} d\eta a(\eta) \cos(k\eta) \chi_k(\eta) \chi_{|\mathbf{k}-\mathbf{p}|}(\eta) \right|^2 + \left| \cos(k\eta) \leftrightarrow \sin(k\eta) \right|^2 \right\}. \quad (20)$$

³ We are considering here classical gravitational waves generated by the motion of matter and not vacuum fluctuations amplified by the cosmic expansion.

Here $(\hat{\mathbf{k}}, \hat{\mathbf{p}})$ is the angle between the unit vectors and the factors $\sin(k\eta)$ and $\cos(k\eta)$ come from the harmonic oscillator's Green functions. Taking into account the expansion of the universe from η_{f} , the present day energy density spectrum of the gravitational waves becomes

$$\Omega_{\text{gw}} h^2 = \frac{S_k(\eta_{\text{f}})}{a_{\text{osc}}^4 \rho_{\text{osc}}} \left(\frac{g_0}{g_{\text{osc}}} \right)^{\frac{1}{3}} \Omega_{\text{r0}} h^2, \quad (21)$$

where we have assumed radiation domination, $r_{\text{osc}} \ll 1$, throughout the generation of gravitational waves for simplicity, g refers to the number of relativistic species and $\Omega_{\text{r0}} h^2$ is the energy density of radiation today.

According to equation (11) the typical physical momenta amplified during preheating are $k_* \sim m_\sigma q^{1/4}$ and the gravitational wave signal will also be peaked around k_* . The corresponding frequency today is [15]

$$f_k \sim \frac{k_*}{\rho_{\text{p}}^{1/4}} 10^{10} \text{ Hz} \sim g^{1/2} r_{\text{osc}}^{1/4} a_{\text{p}}^{3w/4} 10^{10} \text{ Hz} \quad (22)$$

where the subscript "p" refers to the time when the peak was formed and w is the equation of state parameter, $w = 1/3$ for $r_{\text{osc}} \ll 1$ and $w = 0$ for $r_{\text{osc}} \sim 1$. The scale factor at the beginning of oscillations has been normalized to unity as before. Using equation (4), which holds for models where the curvaton value during inflation is given by (3), and taking into account the constraint $g\sigma_* \gg H_*$ required to keep χ massive, one finds the bound

$$f_k \gtrsim \sqrt{\frac{H_*}{M_P}} 10^{10} \text{ Hz} . \quad (23)$$

The frequencies accessible to direct detection experiments range from $f \sim 10^{-4} \text{ Hz}$ (LISA) to $f \sim 10^3 \text{ Hz}$ (LIGO). According to equation (23), the gravitational waves produced in the curvaton preheating can fit this observable window only if the inflationary scale is low enough. If the curvaton would decay perturbatively and if there are no significant changes in its effective mass after inflation, the inflationary scale is constrained by $H_* \gtrsim 10^7 \text{ GeV}$ [21]. This would yield $f_k \gtrsim 10^4 \text{ Hz}$ which unfortunately slightly escapes the observable window. However, equation (23) is a rather crude estimate and a proper numerical computation is required to really find out the actual frequency of the gravitational waves. Moreover, the bound $H_* \gtrsim 10^7 \text{ GeV}$ does not directly apply here since the effective curvaton mass increases during the non-linear stages of preheating and the highly non-thermal epoch after the end of preheating may also significantly affect the expansion history of the universe as compared to the perturbative curvaton decay.

Having discussed the frequencies of the gravitational waves, we are now ready to return to the question of their amplitude. Concentrating first on the linear regime where the backreaction of χ particles can be neglected, we may use equation (20). Gravity waves will only be produced when the frequency of the χ modes varies non-adiabatically, near the zeros of the oscillatory curvaton. Near those zeros the equation of motion for the χ field is given by (9) (remember $X = a^{3/2} \chi$) and its solutions are parabolic cylinder functions $W(-\kappa_n^2/2; \pm\sqrt{2}\tau_n)$

$$X_k^n = A_k^n W(-\kappa_n^2/2; \sqrt{2}\tau_n) + B_k^n W(-\kappa_n^2/2; -\sqrt{2}\tau_n) . \quad (24)$$

This may be matched to the adiabatic solutions (8) in the region where the two solutions overlap. Assuming that the occupation numbers (10) always grow with the fastest possible rate, this yields up to a phase factor [15]

$$X_k^n = \sqrt{\frac{2^{1/2}}{\xi_k} \frac{\beta_k^n}{\sqrt{k_*}}} W(-\kappa_n^2/2; -\sqrt{2}\tau_n) , \quad (25)$$

where $\xi_k \equiv \sqrt{1 + e^{-\pi\kappa^2}} - e^{-\pi\kappa^2/2}$. Using this form and switching to the comoving time in (20) we find

$$S_k(\tau_{\text{f}}) = \frac{m_\sigma^6 G}{8\pi^3 q a^6(\tau_{\text{f}})} K^3 \int_{-1}^1 du (1 - u^2)^2 \int dP P^6 \xi_p \xi_{|\mathbf{k}-\mathbf{p}|} n_p(\tau_{\text{f}}) n_{|\mathbf{k}-\mathbf{p}|}(\tau_{\text{f}}) (I_c^2 + I_s^2) , \quad (26)$$

where $K = k/m_\sigma$ and I_s, I_c are integrals of the parabolic cylinder functions. Except for the prefactors in front of the integral and the exact behaviour of the occupation numbers, this result coincides with equation (78) in [15] derived for the gravitational waves produced in the linear regime of preheating after $\lambda\phi^4$ inflation.

Ignoring the differences in n_k for a while, we may then relate the amplitude of the gravitational waves (21) obtained in [15], $(\Omega_{\text{gw}} h^2)_{\lambda\phi^4}$, to the amplitude in our case as

$$\Omega_{\text{gw}} h^2 \sim 10^{27} \left(\frac{m_\sigma}{M_P} \right)^4 (\Omega_{\text{gw}} h^2)_{\lambda\phi^4} , \quad (27)$$

where subleading contributions coming from different values (and powers) of the scale factor and q have not been included. For a curvaton mass at the TeV scale, the amplitude in (27) is suppressed at least by a factor $\sim 10^{-33}$ or so when compared to $(\Omega_{\text{gw}} h^2)_{\lambda\phi^4}$. From figure 5 of [15] one finds $(\Omega_{\text{gw}} h^2)_{\lambda\phi^4} \lesssim 10^{-13}$ for the linear regime in $\lambda\phi^4$ preheating. Using this in (27), we see that the amplitude is hopelessly below the lower limit of observational sensitivity $\Omega_{\text{gw}} h^2 \sim 10^{-18}$ (BBO). This conclusion holds even when differences in the occupation numbers between our case and the $\lambda\phi^4$ model are taken into account. Assuming that both models would exhibit similar effective growth rates μ but the number of oscillations would be larger by a factor ΔN in the curvaton case, the occupations numbers in the curvaton preheating would be enhanced by a factor $\exp(4\pi\mu\Delta N)$ as compared to the $\lambda\phi^4$ case. This would enhance the amplitude (27) by $\exp(8\pi\mu\Delta N)$ but it would still be difficult to compensate for the suppression $(m_\sigma/M_P)^4$ in (27). For example, for the case $\mu \simeq 0.07$ shown in Fig. (1) one would need $\Delta N \gtrsim 37$, which is more than the number of oscillations taking place during the linear stage, to climb above the threshold $\Omega_{\text{gw}} h^2 \sim 10^{-18}$. We therefore expect that no directly observable gravitational waves will be produced in the *linear* regime of the curvaton preheating.

However, the non-linear stages that follow the linear part of the resonance can give rise to significant production of gravitational waves and the dominant contribution to the total energy density of gravitational waves is typically generated during this stage [14, 15]. Moreover, the non-linear stages are particularly important in the curvaton model since the backreaction effects always become important at the final stages of preheating as discussed above. According to [15], the gravitational wave amplitude from these stages as observed today can be estimated as

$$\Omega_{\text{gw}} h^2 \sim 10^{-6} (R_p H_p)^2, \quad (28)$$

where R_p is the characteristic physical scale where parametric resonance is efficient and H_p is the expansion rate at the time when it occurs. Setting $R_p \sim 1/k_*$ and using equations (3) and (4) we find

$$\Omega_{\text{gw}} h^2 \sim \frac{10^{-6}}{g x^2} \left(\frac{m_\sigma}{H_*} \right)^3 \lesssim \frac{10^{-6}}{x^2} \frac{m_\sigma}{H_*} \quad (29)$$

where we have defined $x \equiv m/H_p$ and also used the condition $g\sigma_* \gg H_*$ in the last step. The parameter x is proportional to the number of oscillation cycles and contributes a factor of few tens. The amplitude of the gravitational waves is therefore typically large enough to be in the detectable range, provided that the mass of the curvaton is not vastly smaller than the inflationary scale H_* . Curvaton models with a sufficiently low inflationary scale, required to fulfill the frequency condition (23), might therefore produce a detectable signal of stochastic gravitational waves. This is an interesting outcome in light of supersymmetric models [22], for example, where the natural curvaton mass would be around the TeV scale. If the inflationary scale is not too much higher, these models could yield directly observable gravitational waves. However, in order to really confirm the qualitative outcome as well as to find out the exact relation between properties of the gravitational waves and the curvaton mass and inflationary scale, one would need to resort to a numerical study.

DISCUSSION

Up to now, discussions of curvaton dynamics have mostly focused on perturbative decay of the curvaton condensate. Here we point out that if the curvaton is about to decay directly into bosonic degrees of freedom, it is highly likely that there will be a non-perturbative preheating stage before the perturbative decay and reheating. The curvaton preheating is qualitatively similar to the conventional inflationary preheating although minor differences can arise due to the fact that the curvaton does not need to dominate the universe during the preheating stage. We have also found that the end of the curvaton preheating is always dominated by the effects of backreaction of the amplified light boson fields. This is different from the inflationary case, where the importance of the backreaction depends on the parameters. Although we have not discussed the details of backreaction, we expect to find no significant differences as compared to the inflationary case here either. This is because the possible differences between the curvaton preheating and the inflationary preheating stem from differences in the expansion of the universe which are unimportant for the final stages of preheating. However, to complete the analysis, one should study numerically the thermalization of the universe after the end of preheating. Since the curvature perturbations generated in the curvaton scenario are sensitive to the evolution history before the curvaton decay, these stages might potentially affect even the gaussian perturbations and therefore yield interesting constraints on the curvaton model itself.

Numerics is also needed to resolve the issue of stochastic gravitational waves. Based on simple estimates we have postulated that models with a sufficiently low inflationary scale could produce gravitational waves with directly detectable frequencies and amplitudes. For example, supersymmetric models [22] with the curvaton mass around the

TeV scale could give rise to observable gravitational waves provided that the inflationary scale is not several orders of magnitude higher. However, our results are still preliminary and a full numerical simulation, subject to a future research, is required before any firm conclusions can be drawn.

Likewise, the possible enhancement of the non-gaussianities in curvaton preheating is best addressed by numerical simulations. Since curvaton preheating is very similar to the inflaton preheating, we may expect enhancements similar as has been found numerically by Chambers and Rajantie [13]. It therefore seems that curvaton decay by a parametric resonance, in addition to being a natural stage in the curvaton scenario, could also have interesting observational consequences.

Acknowledgements

We wish to thank J. F. Dufaux for critical comments as well as A. Rajantie and A. Tranberg for illuminating discussions. This work was supported by the EU 6th Framework Marie Curie Research and Training network "UniverseNet" (MRTN-CT-2006-035863) and partly by Academy of Finland grant 114419. S.N. is supported by the GRASPANP Graduate School.

-
- [1] K. Enqvist and M. S. Sloth, Nucl. Phys. B **626** (2002) 395 [arXiv:hep-ph/0109214].
 - [2] D. H. Lyth and D. Wands, Phys. Lett. B **524** (2002) 5 [arXiv:hep-ph/0110002].
 - [3] T. Moroi and T. Takahashi, Phys. Lett. B **522** (2001) 215 [Erratum-ibid. B **539** (2002) 303] [arXiv:hep-ph/0110096].
 - [4] A. D. Linde and V. F. Mukhanov, Phys. Rev. D **56** (1997) 535 [arXiv:astro-ph/9610219];
S. Mollerach, Phys. Rev. D **42** (1990) 313.
 - [5] N. Bartolo and A. R. Liddle, Phys. Rev. D **65** (2002) 121301 [arXiv:astro-ph/0203076];
S. Gupta, K. A. Malik and D. Wands, Phys. Rev. D **69** (2004) 063513 [arXiv:astro-ph/0311562];
F. Ferrer, S. Rasanen and J. Valiviita, JCAP **0410** (2004) 010 [arXiv:astro-ph/0407300].
 - [6] D. H. Lyth, C. Ungarelli and D. Wands, Phys. Rev. D **67**, 023503 (2003) [arXiv:astro-ph/0208055].
 - [7] N. Bartolo, S. Matarrese and A. Riotto, Phys. Rev. D **69**, 043503 (2004) [arXiv:hep-ph/0309033];
C. Gordon and K. A. Malik, Phys. Rev. D **69**, 063508 (2004) [arXiv:astro-ph/0311102];
K. A. Malik and D. H. Lyth, JCAP **0609**, 008 (2006) [arXiv:astro-ph/0604387];
M. Sasaki, J. Valiviita and D. Wands, Phys. Rev. D **74**, 103003 (2006) [arXiv:astro-ph/0607627];
J. Valiviita, M. Sasaki and D. Wands, [arXiv:astro-ph/0610001];
H. Assadullahi, J. Valiviita and D. Wands, Phys. Rev. D **76** (2007) 103003 [arXiv:0708.0223 [hep-ph]].
 - [8] K. Enqvist and S. Nurmi, JCAP **0510** (2005) 013 [arXiv:astro-ph/0508573].
 - [9] N. Bartolo, S. Matarrese, A. Riotto and A. Vaihkonen, Phys. Rev. D **76** (2007) 061302 [arXiv:0705.4240 [astro-ph]].
 - [10] L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. Lett. **73** (1994) 3195 [arXiv:hep-th/9405187];
Y. Shtanov, J. H. Traschen and R. H. Brandenberger, Phys. Rev. D **51** (1995) 5438 [arXiv:hep-ph/9407247];
J. H. Traschen and R. H. Brandenberger, Phys. Rev. D **42** (1990) 2491.
 - [11] L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. D **56** (1997) 3258 [arXiv:hep-ph/9704452].
 - [12] K. Enqvist, A. Jokinen, A. Mazumdar, T. Multamaki and A. Vaihkonen, Phys. Rev. Lett. **94** (2005) 161301 [arXiv:astro-ph/0411394];
A. Jokinen and A. Mazumdar, JCAP **0604** (2006) 003 [arXiv:astro-ph/0512368].
 - [13] A. Chambers and A. Rajantie, Phys. Rev. Lett. **100** (2008) 041302 [arXiv:0710.4133 [astro-ph]];
A. Chambers and A. Rajantie, arXiv:0805.4795 [astro-ph].
 - [14] S. Y. Khlebnikov and I. I. Tkachev, Phys. Rev. D **56** (1997) 653 [arXiv:hep-ph/9701423];
R. Easther and E. A. Lim, JCAP **0604** (2006) 010 [arXiv:astro-ph/0601617];
J. Garcia-Bellido and D. G. Figueroa, Phys. Rev. Lett. **98** (2007) 061302 [arXiv:astro-ph/0701014];
J. Garcia-Bellido, D. G. Figueroa and A. Sastre, Phys. Rev. D **77** (2008) 043517 [arXiv:0707.0839 [hep-ph]].
 - [15] J. F. Dufaux, A. Bergman, G. N. Felder, L. Kofman and J. P. Uzan, Phys. Rev. D **76** (2007) 123517 [arXiv:0707.0875 [astro-ph]].
 - [16] B. Abbott *et al.* [LIGO Collaboration], Astrophys. J. **659** (2007) 918 [arXiv:astro-ph/0608606]; LIGO Home Page: <http://www.ligo.caltech.edu/>.
 - [17] S. A. Hughes, arXiv:0711.0188 [gr-qc]; LISA Home Page: <http://lisa.esa.int>.
 - [18] V. Corbin and N. J. Cornish, Class. Quant. Grav. **23** (2006) 2435 [arXiv:gr-qc/0512039];
G. M. Harry, P. Fritschel, D. A. Shaddock, W. Folkner and E. S. Phinney, Class. Quant. Grav. **23** (2006) 4887 [Erratum-ibid. **23** (2006) 7361];
BBO Home Page: <http://universe.nasa.gov/new/program/bbo.html>.
 - [19] M. Postma and A. Mazumdar, JCAP **0401** (2004) 005 [arXiv:hep-ph/0304246].
 - [20] K. Dimopoulos, G. Lazarides, D. Lyth and R. Ruiz de Austri, Phys. Rev. D **68** (2003) 123515 [arXiv:hep-ph/0308015].

- [21] D. H. Lyth, “Can the curvaton paradigm accommodate a low inflation scale,” *Phys. Lett. B* **579** (2004) 239 [arXiv:hep-th/0308110].
 K. Dimopoulos, D. H. Lyth and Y. Rodriguez, “Low scale inflation and the curvaton mechanism,” *JHEP* **0502** (2005) 055 [arXiv:hep-ph/0411119].
- [22] R. Allahverdi, K. Enqvist, A. Jokinen and A. Mazumdar, *JCAP* **0610** (2006) 007 [arXiv:hep-ph/0603255];
 K. Enqvist, S. Kasuya and A. Mazumdar, *Phys. Rev. Lett.* **90**, 091302 (2003) [arXiv:hep-ph/0211147];
 K. Enqvist, A. Jokinen, S. Kasuya and A. Mazumdar, *Phys. Rev. D* **68**, 103507 (2003) [arXiv:hep-ph/0303165];
 K. Enqvist, S. Kasuya and A. Mazumdar, *Phys. Rev. Lett.* **93**, 061301 (2004) [arXiv:hep-ph/0311224];
 K. Enqvist, A. Mazumdar and A. Perez-Lorenzana, *Phys. Rev. D* **70**, 103508 (2004) [arXiv:hep-th/0403044];
 K. Enqvist, A. Mazumdar and M. Postma, *Phys. Rev. D* **67**, 121303 (2003) [arXiv:astro-ph/0304187];
 M. Postma, *Phys. Rev. D* **67**, 063518 (2003) [arXiv:hep-ph/0212005];
 S. Kasuya, M. Kawasaki and F. Takahashi, *Phys. Lett. B* **578**, 259 (2004) [arXiv:hep-ph/0305134];
 A. Mazumdar and M. Postma, *Phys. Lett. B* **573** (2003) 5 [Erratum-ibid. *B* **585** (2004) 295] [arXiv:astro-ph/0306509];
 R. Allahverdi, *Phys. Rev. D* **70** (2004) 043507 [arXiv:astro-ph/0403351];
 M. Ikegami and T. Moroi, *Phys. Rev. D* **70**, 083515 (2004) [arXiv:hep-ph/0404253];
 A. Mazumdar, *Phys. Rev. Lett.* **92**, 241301 (2004) [arXiv:hep-ph/0306026];
 A. Mazumdar and A. Perez-Lorenzana, *Phys. Rev. Lett.* **92**, 251301 (2004) [arXiv:hep-ph/0311106];
 A. Mazumdar and A. Perez-Lorenzana, *Phys. Rev. D* **70**, 083526 (2004) [arXiv:hep-ph/0406154].